

PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

2014 Recommendations for Pediatric Preventive Health Care
COMMITTEE ON PRACTICE AND AMBULATORY MEDICINE, BRIGHT
FUTURES PERIODICITY SCHEDULE WORKGROUP
Pediatrics; originally published online February 24, 2014;
DOI: 10.1542/peds.2013-4096

The online version of this article, along with updated information and services, is
located on the World Wide Web at:
<http://pediatrics.aappublications.org/content/early/2014/02/18/peds.2013-4096.citation>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2014 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™





POLICY STATEMENT

2014 Recommendations for Pediatric Preventive Health Care

COMMITTEE ON PRACTICE AND AMBULATORY MEDICINE, 2012–2013

Geoffrey R. Simon, MD, FAAP, Chairperson
Cynthia Baker, MD, FAAP
Graham A. Barden, III, MD, FAAP
Oscar W. Brown, MD, FAAP
Amy Hardin, MD, FAAP
Herschel R. Lessin, MD, FAAP
Kelley Meade, MD, FAAP
Scot Moore, MD, FAAP
Chadwick T. Rodgers, MD, FAAP

FORMER COMMITTEE MEMBERS

Lawrence D. Hammer, MD, FAAP, Chairperson
Edward S. Curry, MD, FAAP*
James J. Laughlin, MD, FAAP

STAFF

Elizabeth Sobczyk, MPH, MSW

BRIGHT FUTURES PERIODICITY SCHEDULE WORKGROUP

Edward S. Curry, MD, FAAP*
Paula M. Duncan, MD, FAAP
Joseph F. Hagan, Jr, MD, FAAP
Alex R. Kemper, MD, MPH, MS, FAAP
Judith S. Shaw, EdD, MPH, RN, FAAP
Jack T. Swanson, MD, FAAP

STAFF

Jane B. Bassewitz, MA

COMMITTEE ON PRACTICE AND AMBULATORY MEDICINE, BRIGHT FUTURES PERIODICITY SCHEDULE WORKGROUP

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

The guidance in this statement does not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

All policy statements from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

www.pediatrics.org/cgi/doi/10.1542/peds.2013-4096

doi:10.1542/peds.2013-4096

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2014 by the American Academy of Pediatrics

FREE

*Dr Curry serves as the Committee on Practice and Ambulatory Medicine liaison to Bright Futures and is a member of the Bright Futures Steering Committee.

Bright Futures/American Academy of Pediatrics



Developmental, psychosocial, and chronic disease issues for children and adolescents may require frequent counseling and treatment visits separate from preventive care visits.

Refer to the specific guidance by age as listed in *Bright Futures* guidelines (Hagan JF, Shaw JS, Duncan PM, eds. *Bright Futures Guidelines for Health Supervision of Infants, Children and Adolescents*. 3rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2008).

Copyright © 2014 by the American Academy of Pediatrics.

No part of this statement may be reproduced in any form or by any means without prior written permission from the American Academy of Pediatrics except for one copy for personal use.

[illegible]

1. If a child comes under care for the first time at any point on the schedule, or if any items are not accomplished at the suggested age, the schedule should be brought up to date at the earliest possible time.
2. A prenatal visit is recommended for parents who are at high risk, for first-time parents, and for those who request a conference. The prenatal visit should include anticipatory guidance, pertinent medical history, and a discussion of benefits of breastfeeding and planned method of feeding, per the 2009 AAP statement "The Prenatal Visit" (<http://pediatrics.aappublications.org/content/124/4/1227.full>).
3. Every infant should have a newborn evaluation after birth, and breastfeeding should be encouraged (and instruction and support should be offered).
4. Every infant should have an evaluation within 3 to 5 days of birth and within 48 to 72 hours after discharge from the hospital to include evaluation for feeding and jaundice. Breastfeeding infants should receive formal breastfeeding evaluation, and their mothers should receive encouragement and instruction, as recommended in the 2012 AAP statement "Breastfeeding and the Use of Human Milk" (<http://pediatrics.aappublications.org/content/129/3/6827.full>). Newborn infants discharged less than 48 hours after delivery must be examined within 48 hours of discharge, per the 2010 AAP statement "Hospital Stay for Healthy Term Newborns" (<http://pediatrics.aappublications.org/content/125/2/405.full>).
5. Screen, per the 2007 AAP statement "Expert Committee Recommendations Regarding the Prevention, Assessment, and Treatment of Child and Adolescent Overweight and Obesity: Summary Report" (http://pediatrics.aappublications.org/content/120/Supplement_4/S164.full).
6. Blood pressure measurement in infants and children with specific risk conditions should be performed at visits before age 3 years.
7. If the patient is uncooperative, rescreen within 6 months, per the 2007 AAP statement "Eye Examination in Infants, Children, and Young Adults by Pediatricians" (<http://pediatrics.aappublications.org/content/111/4/902.abstract>).
8. All newborns should be screened, per the AAP statement "Year 2007 Position Statement: Principles and Guidelines for Early Hearing Detection and Intervention Programs" (<http://pediatrics.aappublications.org/content/120/4/898.full>).
9. See 2006 AAP statement "Identifying Infants and Young Children With Developmental Disorders in the Medical Home: An Algorithm for Developmental Surveillance and Screening" (<http://pediatrics.aappublications.org/content/118/1/405.full>).
10. Screening should occur per the 2007 AAP statement "Identification and Evaluation of Children with Autism Spectrum Disorders" (<http://pediatrics.aappublications.org/content/120/5/1183.full>).
11. A recommended screening tool is available at <http://www.ceasars-boston.org/CRAFT/index.php>.
12. Recommended screening using the Patient Health Questionnaire (PHQ)-2 or other tools available in the GLAD-PC toolkit and at http://www.aap.org/en-us/advocacy-and-policy/aap-health-initiatives/Mental-Health/Documents/MH_ScreeningChart.pdf.
13. At each visit, age-appropriate physical examination is essential, with infant totally unclothed and older children undressed and suitably draped. See 2011 AAP statement "Use of Chaperones During the Physical Examination of the Pediatric Patient" (<http://pediatrics.aappublications.org/content/127/5/991.full>).
14. These may be modified, depending on entry point into schedule and individual need.
15. The Recommended Uniform Newborn Screening Panel (<http://www.hrsa.gov/advisorycommittees/mchadv/ns/heritabledisorders/recommendedpanel/uniformscreeningpanel.pdf>), as determined by The Secretary's Advisory Committee on Heritable Disorders in Newborns and Children, and state newborn screening laws/regulations (<http://genes-c-us.uthscsa.edu/sites/genes-c-us/files/nbstdisorders.pdf>), establish the criteria for and coverage of newborn screening procedures and programs. Follow-up must be provided, as appropriate, by the pediatrician.
16. Screening for critical congenital heart disease using pulse oximetry should be performed in newborns, after 24 hours of age, before discharge from the hospital, per the 2011 AAP statement "Endorsement of Health and Human Services Recommendation for Pulse Oximetry Screening for Critical Congenital Heart Disease" (<http://pediatrics.aappublications.org/content/129/1/190.full>).
17. Schedules, per the AAP Committee on Infectious Diseases, are available at: <http://aapredbook.aappublications.org/site/resources/zschedules.xhtml>. Every visit should be an opportunity to update and complete a child's immunizations.
18. See 2010 AAP statement "Diagnosis and Prevention of Iron Deficiency and Iron Deficiency Anemia in Infants and Young Children (0-3 Years of Age)" (<http://pediatrics.aappublications.org/content/126/5/1040.full>).
19. For children at risk of lead exposure, see the 2012 CDC Advisory Committee on Childhood Lead Poisoning Prevention statement "Low Level Lead Exposure Harms Children: A Renewed Call for Primary Prevention" (http://www.cdc.gov/nceh/lead/ACCLPP/Final_Document_030712.pdf).
20. Perform risk assessments or screenings as appropriate, based on universal screening requirements for patients with Medicaid or in high prevalence areas.
21. Tuberculosis testing per recommendations of the Committee on Infectious Diseases, published in the current edition of *Red Book: Report of the Committee on Infectious Diseases*. Testing should be performed on occurrence of high-risk factors.
22. See AAP-endorsed 2011 guidelines from the National Heart Blood and Lung Institute, "Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents" (http://www.nhlbi.nih.gov/guidelines/cvd_ped/index.htm).
23. Adolescents should be screened for sexually transmitted infections (STIs) per recommendations in the current edition of the *AAP Red Book: Report of the Committee on Infectious Diseases*. Additionally, all adolescents should be screened for HIV according to the AAP statement (<http://pediatrics.aappublications.org/content/128/5/1023.full>) once between the ages of 16 and 18, making every effort to preserve confidentiality of the adolescent. Those at increased risk of HIV infection, including those who are sexually active, participate in injection drug use, or are being tested for other STIs, should be tested for HIV and reassessed annually.
24. See USPSTF recommendations (<http://www.uspreventiveservicestaskforce.org/uspstf/uspsscerv.htm>). Indications for pelvic examinations prior to age 21 are noted in the 2010 AAP statement "Gynecologic Examination for Adolescents in the Pediatric Office Setting" (<http://pediatrics.aappublications.org/content/126/3/583.full>).
25. Refer to a dental home, if available. If not available, perform a risk assessment (<http://www2.aap.org/oralhealth/docs/RiskAssessmentTool.pdf>). If primary water source is deficient in fluoride, consider oral fluoride supplementation. For those at high risk, consider application of fluoride varnish for caries prevention. See 2008 AAP statement "Preventive Oral Health Intervention for Pediatricians" (<http://pediatrics.aappublications.org/content/122/6/1387.full>) and 2009 AAP statement "Oral Health Risk Assessment Timing and Establishment of the Dental Home" (<http://pediatrics.aappublications.org/content/111/5/1113.full>).

KEY ● = to be performed ★ = risk assessment to be performed with appropriate action to follow, if positive ◀ ● ▶ = range during which a service may be provided

Summary of changes made to the 2014 AAP Recommendations for Preventive Pediatric Health Care (Periodicity Schedule)

Changes to Developmental/Behavioral Assessment

- **Alcohol and Drug Use Assessment-** Information regarding a recommended screening tool (CRAFTT) was added.
- **Depression-** Screening for depression at ages 11 through 21 has been added, along with suggested screening tools.

Changes to Procedures

- **Dyslipidemia screening-** An additional screening between 9 and 11 years of age has been added. The reference has been updated to the AAP-endorsed National Heart Blood and Lung Institute policy (http://www.nhlbi.nih.gov/guidelines/cvd_ped/index.htm).
- **Hematocrit or hemoglobin-** A risk assessment has been added at 15 and 30 months. The reference has been updated to the current AAP policy (<http://pediatrics.aappublications.org/content/126/5/1040.full>).
- **STI/HIV screening-** A screen for HIV has been added between 16 and 18 years. Information on screening adolescents for HIV has been added in the footnotes. STI screening now references recommendations made in the AAP Red Book. This category was previously titled "STI Screening."
- **Cervical dysplasia-** Adolescents should no longer be routinely screened for cervical dysplasia until age 21. Indications for pelvic exams prior to age 21 are noted in the 2010 AAP statement "Gynecologic Examination for Adolescents in the Pediatric Office Setting" (<http://pediatrics.aappublications.org/content/126/3/583.full>).
- **Critical Congenital Heart Disease-** Screening for critical congenital heart disease using pulse oximetry should be performed in newborns, after 24 hours of age, before discharge from the hospital, per the 2011 AAP statement, "Endorsement of Health and Human Services Recommendation for Pulse Oximetry Screening for Critical Congenital Heart Disease" (<http://pediatrics.aappublications.org/content/129/1/190.full>).

For several recommendations, the AAP Policy has been updated since 2007 but there have been no changes in the timing of recommendations on the Periodicity Schedule. These include:

- Footnote 2- The Prenatal Visit (2009): <http://pediatrics.aappublications.org/content/124/4/1227.full>
- Footnote 4- Breastfeeding and the Use of Human Milk (2012): <http://pediatrics.aappublications.org/content/129/3/e827.full> and Hospital Stay for Healthy Term Newborns (2010): <http://pediatrics.aappublications.org/content/125/2/405.full>
- Footnote 8- Year 2007 Position Statement: Principles and Guidelines for Early Hearing Detection and Intervention Programs (2007): <http://pediatrics.aappublications.org/content/120/4/898.full>
- Footnote 10- Identification and Evaluation of Children with Autism Spectrum Disorders (2007): <http://pediatrics.aappublications.org/content/120/5/1183.full>
- Footnote 17- Immunization Schedules (2013): <http://aapredbook.aappublications.org/site/resources/IZSchedule0-6yrs.pdf>, <http://aapredbook.aappublications.org/site/resources/IZSchedule7-18yrs.pdf>, and <http://aapredbook.aappublications.org/site/resources/IZScheduleCatchup.pdf>
- Footnote 19- CDC Advisory Committee on Childhood Lead Poisoning Prevention statement "Low Level Lead Exposure Harms Children: A Renewed Call for Primary Prevention" (2012): http://www.cdc.gov/nceh/lead/ACCLPP/Final_Document_030712.pdf
- Footnote 22- AAP-endorsed guideline "Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents" (2011): http://www.nhlbi.nih.gov/guidelines/cvd_ped/index.htm
- Footnote 25- Preventive Oral Health Intervention for Pediatricians (2008): <http://pediatrics.aappublications.org/content/122/6/1387.full> and Oral Health Risk Assessment Timing and Establishment of the Dental Home (2009): <http://pediatrics.aappublications.org/content/111/5/1113.full>. Additional information from the policies regarding fluoride supplementation and fluoride varnish has been added to the footnote.

New references were added for several footnotes, also with no change to recommendations in the Periodicity Schedule:

- Footnote 5- Expert Committee Recommendations Regarding the Prevention, Assessment, and Treatment of Child and Adolescent Overweight and Obesity: Summary Report (2007): http://pediatrics.aappublications.org/content/120/Supplement_4/S164.full
- Footnote 13- Use of Chaperones During the Physical Examination of the Pediatric Patient (2011): <http://pediatrics.aappublications.org/content/127/5/991.full>
- Footnote 15- The Recommended Uniform Newborn Screening Panel (<http://www.hrsa.gov/advisorycommittees/mchbadvisory/heritabledisorders/recommendedpanel/uniformscreeningpanel.pdf>), as determined by The Secretary's Advisory Committee on Heritable Disorders in Newborns and Children, and state newborn screening laws/regulations (<http://genes-r-us.uthscsa.edu/sites/genes-r-us/files/nbsdisorders.pdf>), establish the criteria for and coverage of newborn screening procedures and programs. Follow-up must be provided, as appropriate, by the pediatrician.

For consistency, the title of "Tuberculin Test" has been changed to "Tuberculosis Testing." The title of "Newborn Metabolic/Hemoglobin Screening" has been changed to "Newborn Blood Screening."

2014 Recommendations for Pediatric Preventive Health Care
COMMITTEE ON PRACTICE AND AMBULATORY MEDICINE, BRIGHT
FUTURES PERIODICITY SCHEDULE WORKGROUP
Pediatrics; originally published online February 24, 2014;
DOI: 10.1542/peds.2013-4096

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/early/2014/02/18/peds.2013-4096.citation>

Citations

This article has been cited by 1 HighWire-hosted articles:
<http://pediatrics.aappublications.org/content/early/2014/02/18/peds.2013-4096.citation#related-urls>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://pediatrics.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://pediatrics.aappublications.org/site/misc/reprints.xhtml>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2014 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

